

The Joint Commission 10 Harmonized Stroke Measures
Facilities that are Primary Stroke Center (PSC) Certified and/or those considering PSC Certification should strive for $\geq 80\%$ compliance with these 10 measures

DVT Prophylaxis Stroke-1

Percent of patients with an ischemic stroke or a hemorrhagic stroke and who are non-ambulatory who receive DVT prophylaxis by end of hospital day two.

Antithrombotics at Discharge Stroke-2

Percent of patients with an ischemic stroke prescribed antithrombotic therapy at discharge.

Anticoagulation for Atrial Fibrillation/Flutter Stroke-3

Percent of patients with an ischemic stroke with atrial fibrillation/flutter receiving anticoagulation therapy.

t-PA Administration Stroke-4

Percent of acute ischemic stroke patients who arrive at the hospital within 120 minutes (2 hours) of time last known well and for whom IV t-PA was initiated at this hospital within 180 minutes (3 hours) of time last known well.

Early Antithrombotics Stroke-5

Percent of patients with ischemic stroke who receive antithrombotic therapy by the end of hospital day two.

LDL 100 - Discharged on Cholesterol Reducers Stroke-6

Percent of Ischemic stroke patients with LDL > 100, or LDL not measured, or on cholesterol-reducer prior to admission, who are discharged on cholesterol reducing drugs.

*** Dysphagia Screen Stroke-7**

Percent of patients with ischemic or hemorrhagic stroke who undergo screening for dysphagia with an evidence-based bedside testing protocol approved by the hospital before being given any food, fluids, or medication by mouth.

Stroke Education Stroke-8

Percent of patients with ischemic or hemorrhagic stroke or their caregivers who were given education and/or educational materials during the hospital stay addressing ALL of the following: personal risk factors for stroke, warning signs for stroke, activation of emergency medical system, need for follow-up after discharge, and medications prescribed at discharge.

*** Smoking Cessation Stroke-9**

Percent of patients with ischemic or hemorrhagic stroke with a history of smoking cigarettes, who are, or whose caregivers are, given smoking cessation advice or counseling during hospital stay.

Assessed for Rehabilitation Services Stroke-10

Percent of patients with an ischemic or hemorrhagic stroke who were assessed for rehabilitation services.

* CMS/NQF Recommended Stroke Core Measures to exclude Dysphagia and Smoking Cessation Effective October 1, 2009

Source: Joint Commission, American Heart Association, American Stroke Association, CDC-PCNASR (Paul Coverdell National Acute Stroke Registry)

Document prepared by Cristine Small, PCMH, ENCSN QI Workgroup, Chair <http://www.encsn.org/qi.htm>

Revised January 2008, Revised February 2009